


National Library of Medicine

My NC
[\[Sign In\]](#) [\[Register\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for

☒ Limits ☐ Preview/Index ☐ History ☐ Clipboard ☐ Details

Field: **Title**

Display Show Sort by Send to

All: 1 Review: 0

[About Entrez](#)[Text Version](#)[Entrez PubMed](#)[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)[PubMed Services](#)
[Journals Database](#)
[MeSH Database](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[My NCBI \(Cubby\)](#)[Related Resources](#)
[Order Documents](#)
[NLM Catalog](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)☐ 1: Biochimie. 1989 Jul;71(7):857-63.[Related Articles, Links](#)

Integration and germ line transmission of foreign genes microinjected into fertilized trout eggs.

Guyomard R, Chourrout D, Leroux C, Houdebine LM, Pourrain F.

Laboratoire de Genetique des Poissons, Institut National de la Recherche Agronomique, Castanet-Tolosan, France.

Persistence, integration into host genome, germ line transmission and expression of foreign genes microinjected into cytoplasm of fertilized rainbow trout eggs has been examined. Foreign DNA persisted as large random concatamers in approximately 50% of 6 to 12 month-old trout and exhibited a mosaic pattern between tissues. In some cases, free concatamers were observed indicating that extrachromosomal replication occurred in trout. Approximately 50% of the males had the foreign sequences in sperm DNA and all the examined animals transmitted these sequences to their progeny. The percentage of transgenic offsprings ranged from 10 to 30% and putative junction fragments were identified in Southern blot analysis in some of them. These results strongly support the hypothesis that the injected genes became integrated into the genome host, most likely after the first round of chromosomal replication. We also examined the expression of the microinjected plasmids which contained viral or mammalian promoters linked to human or rat growth hormone gene. In no case could exogenous growth hormone be detected.

PMID: 2505865 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)
[NCBI | NLM | NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)